The Paradox of the Fit Heart

We like to believe that a life spent pedalling keeps Father Time firmly at bay. After all, surely all those miles count for something. Every now and then, though, the body quietly reminds us that fitness, enthusiasm and an expensive bike do not grant full immunity. Since joining the club, I have spoken to several members about those strange moments when the heart rate seems to take matters into its own hands, surging as if a hill sprint has been launched despite the pedals barely turning. It can be deeply unsettling, particularly when you are certain you were *not* attacking.

One of the more common rhythm changes is **Atrial Fibrillation**. In simple terms, the heart slips out of its usual steady rhythm and begins to beat irregularly, sometimes more quickly than expected. On the bike this can feel like a fluttering in the chest or throat, breathing that suddenly feels harder than it should, or a sense that the engine will not quite respond even though the legs are willing and keen. For some riders there is no obvious sensation at all, just an unexplained drop in form where familiar efforts feel strangely difficult. The key feature is that it feels different from normal fatigue, as if the pacing has gone awry rather than the power simply running out. In other words, it does not feel like “I’m unfit” so much as “something isn’t quite right here”.

Another common condition is **Supraventricular Tachycardia**. This is defined by a sudden, inappropriate acceleration of the heart rate, often jumping very quickly to something in the region of 150 to 200 beats per minute rather than building gradually. The key features are that the heartbeat becomes much faster than expected for the effort being made and that it is usually regular rather than erratic. That is why riders often notice it immediately and describe it as a switch being flipped, usually at a moment when they were feeling perfectly comfortable a second earlier.

It is also worth remembering that heart rhythm issues do not always reveal themselves clearly on a heart-rate monitor. Sometimes there is a stubbornly high or erratic reading, but just as often the screen looks perfectly normal and entirely unconcerned. For that reason, the numbers are best treated as a guide rather than a verdict. How you feel still matters most. Sensations such as fluttering, breathlessness out of proportion to effort, sudden weakness or light-headedness are worth paying attention to whether the data agrees or not. Used

sensibly, heart-rate monitoring can support good decisions, but it should never override what your body is telling you, however much you like your gadgets.

If an episode starts during a club ride, there are a few simple things that may help it settle. Easing the effort straight away is key. Gentle pedalling or stopping altogether often allows the heart to return to its normal rhythm without further intervention, and nobody will think less of you for soft pedalling home. Deliberately slowing your breathing can also help, focusing on steady, controlled breaths rather than short, shallow ones. Staying calm and avoiding sudden exertion reduces adrenaline release, which can otherwise prolong symptoms.

Taking on fluid little and often can help, particularly if you have been sweating or riding in warm conditions, as dehydration can worsen symptoms. There is no need to force large volumes at once. Small, regular sips are usually sufficient while things settle. Caffeine is not a trigger for most people, but it is reasonable to avoid further stimulants until things have settled and the situation has been properly assessed.

Most importantly, letting someone know how you are feeling removes any pressure to push on heroically and allows the group to support you appropriately. If symptoms persist, worsen, or are accompanied by chest pain, marked breathlessness, dizziness or faintness, stopping the ride and seeking medical help is essential.

Even if the episode settles, it is wise to arrange further assessment. A GP is a good starting point, and referral to a cardiologist is often appropriate. Further tests may include heart rhythm monitoring and imaging, not because something terrible is assumed, but because clarity is reassuring and guesswork is not a medical strategy.

If you need to seek advice, it helps to describe the episode clearly. Note what you were doing at the time, how hard you were exercising, how long the symptoms lasted and how they settled. How it felt is just as important as any numbers, whether that was fluttering, unexpected breathlessness, light-headedness or a sudden loss of power. If you had a sense of whether the rhythm felt irregular or simply very fast and steady, that’s worth mentioning. You can say if you wondered whether it might have been something like AF or SVT, but describing the pattern matters more than getting the name right.

Set a pact with yourself: It’s worth ending on a note of perspective, and a little caution about what you read. AI searches and online summaries often suggest a simple link between older athletes and electrical heart conditions, which can sound more alarming than it needs to be. Regular exercise remains one of the best things we can do for long-term health, benefiting the heart, muscles, metabolism and mental wellbeing in ways no pill can match. The sensible response is not to stop exercising, but to decide in advance that if something feels out of character, you will pause and seek medical advice. What has changed is not that there is a sudden epidemic, but that our understanding of cardiophysiology, and our ability to measure what the heart is doing, has improved dramatically.

Dr Mark Benson